

Product datasheet for TA384313

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Aspartate Aminotransferase (GOT1) Rabbit Monoclonal Antibody [Clone ID: R05-6C1]

Product data:

Product Type: Primary Antibodies

Clone Name: R05-6C1

Applications: IF, IHC, WB

Recommended Dilution: WB: 1/1000

IHC: 1/20-1/100 ICC/IF: 1/100-1/100

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human Aspartate Aminotransferase

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 46 kDa; Observed MW: 46 kDa

Gene Name: glutamic-oxaloacetic transaminase 1

Database Link: Entrez Gene 2805 Human

P17174






Background:


Swiss-Prot Acc.P17174.Biosynthesis of L-glutamate from L-aspartate or L-cysteine. Important regulator of levels of glutamate, the major excitatory neurotransmitter of the vertebrate central nervous system. Acts as a scavenger of glutamate in brain neuroprotection. The aspartate aminotransferase activity is involved in hepatic glucose synthesis during development and in adipocyte glyceroneogenesis. Using L-cysteine as substrate, regulates levels of mercaptopyruvate, an important source of hydrogen sulfide. Mercaptopyruvate is converted into H2S via the action of 3-mercaptopyruvate sulfurtransferase (3MST). Hydrogen sulfide is an important synaptic modulator and neuroprotectant in the brain.

Synonyms: GIG18

Product images:




Western blot analysis of Aspartate Aminotransferase in K562, rat Brain, 3T3, Hela lysates using Aspartate Aminotransferase antibody.



Immunohistochemistry analysis of paraffinembedded Human colon cancer tissue using Aspartate Aminotransferase antibody. Highpressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunocytochemistry analysis of Aspartate Aminotransferase (green) in U87-MG using Aspartate Aminotransferase antibody, and DAPI(blue).