

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA384335S

### Metabotropic Glutamate Receptor 2 (GRM2) Rabbit Monoclonal Antibody [Clone ID: R04-2D1]

#### **Product data:**


| Product Type:           | Primary Antibodies   |
|-------------------------|--|
| Clone Name:             | R04-2D1  |
| Applications:           | IHC, WB  |
| Recommended Dilution:   | WB: 1/2000-1/10000<br>IHC: 1/50  |
| Reactivity:             | Human, Mouse   |
| Host:                   | Rabbit   |
| lsotype:                | lgG  |
| Clonality:              | Monoclonal   |
| Immunogen:              | A synthetic peptide of human Metabotropic Glutamate Receptor 2   |
| Formulation:            | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA  |
| Concentration:          | lot specific   |
| Purification:           | Affinity Purified  |
| Conjugation:            | Unconjugated   |
| Storage:                | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.   |
| Stability:              | 1 year   |
| Predicted Protein Size: | Calculated MW: 99 kDa; Observed MW: 99 kDa   |
| Gene Name:              | glutamate metabotropic receptor 2  |
| Database Link:          | <u>Entrez Gene 2912 Human</u><br><u>Q14416</u>   |
| Background:             | Swiss-Prot Acc.Q14416.G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity. May mediate suppression of neurotransmission or may be involved in synaptogenesis or synaptic stabilization. |
| Synonyms:               | GLUR2; GPRC1B; mGlu2; MGLUR2   |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



## **Product images:**



Western blot analysis of Metabotropic Glutamate Receptor 2 in K562, Hela lysates using Metabotropic Glutamate Receptor 2 antibody.



Immunohistochemistry analysis of paraffinembedded mouse testis using Metabotropic Glutamate Receptor 2 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US