

Product datasheet for TA384339

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GSK3 alpha (GSK3A) Rabbit Monoclonal Antibody [Clone ID: R01-5G6]

Product data:

Product Type: Primary Antibodies

Clone Name: R01-5G6
Applications: IP, WB

Recommended Dilution: WB: 1/1000

IP: 1/20

Reactivity: Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic phosphopeptide corresponding to residues surrounding Tyr216 of human GSK3

alpha (Phosphorylated)

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 51,47 kDa; Observed MW: 47-51 kDa

Gene Name: glycogen synthase kinase 3 alpha

Database Link: P49840




Background:

Swiss-Prot Acc.P49840.Constitutively active protein kinase that acts as a negative regulator in the hormonal control of glucose homeostasis, Wnt signaling and regulation of transcription factors and microtubules, by phosphorylating and inactivating glycogen synthase (GYS1 or GYS2), CTNNB1/beta-catenin, APC and AXIN1. Requires primed phosphorylation of the majority of its substrates. Contributes to insulin regulation of glycogen synthesis by phosphorylating and inhibiting GYS1 activity and hence glycogen synthesis. Regulates glycogen metabolism in liver, but not in muscle. May also mediate the development of insulin resistance by regulating activation of transcription factors. In Wnt signaling, regulates the level and transcriptional activity of nuclear CTNNB1/beta-catenin. Facilitates amyloid precursor protein (APP) processing and the generation of APP-derived amyloid plaques found in Alzheimer disease. May be involved in the regulation of replication in pancreatic beta-cells. Is necessary for the establishment of neuronal polarity and axon outgrowth. Through phosphorylation of the anti-apoptotic protein MCL1, may control cell apoptosis in response to growth factors deprivation.

Synonyms: DKFZp686D0638; OTTHUMP00000198368

Product images:



Western blot detection of Phospho-GSK3 (alpha/beta) (Tyr216/Tyr279) in C6,3T3 cell lysates using Phospho-GSK3 (alpha/beta) (Tyr216/Tyr279) Rabbit mAb(1:1000 diluted).Predicted band size:51,47kDa.Observed band size:47-51kDa.