

Product datasheet for TA384369

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

H2BC19P Rabbit Monoclonal Antibody [Clone ID: R07-8K4]

Product data:

Product Type: Primary Antibodies

Clone Name: R07-8K4

Applications: IHC, IP, WB

Recommended Dilution: WB: 1/2000

IHC: 1/200 IP: 1/20

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human Histone H2B

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 14 kDa; Observed MW: 14 kDa

Gene Name: histone cluster 2, H2bd (pseudogene)

Database Link: Entrez Gene 337874 Human

060814






Background:


Swiss-Prot Acc.O60814.Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Has broad antibacterial activity. May contribute to the formation of the functional antimicrobial barrier of the colonic epithelium, and to the bactericidal activity of amniotic fluid.

Synonyms: HIST2H2BD

Product images:




Western blot analysis of Histone H2B Receptor in 3T3 lysates using Histone H2B antibody.



Western blot analysis of Histone H2B in K562, C6 lysates using Histone H2B antibody.





Immunohistochemistry analysis of paraffinembedded Human colon cancer using Histone H2B antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.