

Product datasheet for TA384438

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HMGA1 Rabbit Monoclonal Antibody [Clone ID: R01-9D3]

Product data:

Product Type: Primary Antibodies

Clone Name: R01-9D3

Applications: IF, IHC, IP, WB

Recommended Dilution: WB: 1/1000

IHC: 1/200 ICC/IF: 1/100 IP: 1/20

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human HMGA1

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 12 kDa; Observed MW: 18 kDa

Gene Name: high mobility group AT-hook 1

Database Link: Entrez Gene 3159 Human

<u>P17096</u>

Background: Swiss-Prot Acc.P17096.HMG-I/Y bind preferentially to the minor groove of A+T rich regions in

double-stranded DNA. It is suggested that these proteins could function in nucleosome phasing and in the 3'-end processing of mRNA transcripts. They are also involved in the

transcription regulation of genes containing, or in close proximity to A+T-rich

regions.RegionsFeature keyPosition(s)Description







Synonyms:

HMG-I(Y); HMG-R; HMGA1A; HMGIY; MGC4242; MGC4854; MGC12816; OTTHUMP00000016223; OTTHUMP00000016224

Product images:




Western blot analysis of HMGA1 in Hela, C6, MCF-7 lysates using HMGA1 antibody.



Immunohistochemistry analysis of paraffinembedded Human tonsil using HMGA1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunocytochemistry analysis of HMGA1(green) in Hela using HMGA1 antibody,and DAPI(blue)