

Product datasheet for TA384482

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

IFITM3 Rabbit Monoclonal Antibody [Clone ID: R09-615]

Product data:

Product Type: Primary Antibodies

Clone Name: R09-6I5

Applications: IF, IHC, WB **Recommended Dilution:** WB: 1/1000

IHC: 1/50 ICC/IF: 1/50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human Fragilis

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 15 kDa; Observed MW: 15 kDa **Gene Name:** interferon induced transmembrane protein 3

Database Link: Entrez Gene 10410 Human

Q01628






Background:

Swiss-Prot Acc.Q01628.IFN-induced antiviral protein which disrupts intracellular cholesterol homeostasis. Inhibits the entry of viruses to the host cell cytoplasm by preventing viral fusion with cholesterol depleted endosomes. May inactivate new enveloped viruses which buds out of the infected cell, by letting them go out with a cholesterol depleted membrane. Active against multiple viruses, including influenza A virus, SARS coronavirus (SARS-CoV), Marburg virus (MARV) and Ebola virus (EBOV), Dengue virus (DNV), West Nile virus (WNV), human immunodeficiency virus type 1 (HIV-1) and vesicular stomatitis virus (VSV). Can inhibit: influenza virus hemagglutinin protein-mediated viral entry, MARV and EBOV GP1,2-mediated viral entry, SARS-CoV S protein-mediated viral entry and VSV G protein-mediated viral entry. Plays a critical role in the structural stability and function of vacuolar ATPase (v-ATPase). Establishes physical contact with the v-ATPase of endosomes which is critical for proper clathrin localization and is also required for the function of the v-ATPase to lower the pH in phagocytic endosomes thus establishing an antiviral state.

Synonyms:


1-8U; interferon-inducible; IP15

Product images:




Western blot analysis of Fragilis in K562, Hela lysates using Fragilis antibody.





Immunohistochemistry of Fragilis in paraffinembedded Human breast cancer tissue using Fragilis Rabbit mAb at dilution 1/5



Immunocytochemistry of Fragilis(green) in Hela cells using Fragilis Rabbit mAb at dilution 1/50, and DAPI(blue)