

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA384730M

### p38 (MAPK14) Rabbit Monoclonal Antibody [Clone ID: R06-7D3]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	R06-7D3
Applications:	IP, WB
Recommended Dilution:	WB: 1/1000 IP: 1/20
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Monoclonal
Immunogen:	A synthetic peptide of human p38
Formulation:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year
Predicted Protein Size:	Calculated MW:41 kDa,Observed MW:41 kDa
Gene Name:	mitogen-activated protein kinase 14
Database Link:	<u>Entrez Gene 1432 Human</u> <u>Q16539</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **P38 (MAPK14) Rabbit Monoclonal Antibody [Clone ID: R06-7D3] - TA384730MBackground:**The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as<br/>an integration point for multiple biochemical signals, and are involved in a wide variety of<br/>cellular processes such as proliferation, differentiation, transcription regulation and<br/>development. This kinase is activated by various environmental stresses and<br/>proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase<br/>kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1<br/>protein with this kinase. The substrates of this kinase include transcription regulator ATF2,<br/>MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the<br/>roles of this kinase in stress related transcription and cell cycle regulation, as well as in<br/>genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding<br/>distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Synonyms: CSBP; CSBP1; CSBP2; CSPB1; EXIP; Mxi2; p38; p38ALPHA; PRKM14; PRKM15; RK; SAPK2A

## **Product images:**



Western blot detection of p38 in Hela,A549,MCF-7,U251 cell lysates using p38 Rabbit mAb(1:1000 diluted).Predicted band size:41kDa.Observed band size:41kDa.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US