

Product datasheet for TA384831

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Myogenin (MYOG) Mouse Monoclonal Antibody [Clone ID: 3G9-3A9-2B3]

Product data:

Product Type: Primary Antibodies

Clone Name: 3G9-3A9-2B3

Applications: IHC

Recommended Dilution: IHC-p: 1/100-1/500

Reactivity: Human Host: Mouse

Isotype: IgG1, kappa
Clonality: Monoclonal

Immunogen: Synthesized peptide derived from human Myogenin

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Concentration: lot specific

Purification: Affinity Chromatography

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Gene Name: myogenin (myogenic factor 4)

Database Link: Entrez Gene 4656 Human


P15173

Background: Swiss-Prot Acc.P15173.


Synonyms: bHLHc3; Myf-4; MYF4; MYOGENIN



Product images:




Immunohistochemical analysis of paraffinembedded Rhabdomyosarcoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffinembedded Rhabdomyosarcoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





Immunohistochemical analysis of paraffinembedded Rhabdomyosarcoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6. was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffinembedded Rhabdomyosarcoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).