

Product datasheet for **TA384917M**

P4HB Rabbit Monoclonal Antibody [Clone ID: R09-1F4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	R09-1F4
Applications:	IHC, IP, WB
Recommended Dilution:	WB: 1/1000 IHC: 1/20 IP: 1/20
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Immunogen:	A synthetic peptide of human P4HB
Formulation:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year
Predicted Protein Size:	Calculated MW: 57 kDa; Observed MW: 57 kDa
Gene Name:	prolyl 4-hydroxylase subunit beta
Database Link:	Entrez Gene 5034 Human P07237




[View online »](#)

Background:


Swiss-Prot Acc.P07237. This multifunctional protein catalyzes the formation, breakage and rearrangement of disulfide bonds. At the cell surface, seems to act as a reductase that cleaves disulfide bonds of proteins attached to the cell. May therefore cause structural modifications of exofacial proteins. Inside the cell, seems to form/rearrange disulfide bonds of nascent proteins. At high concentrations, functions as a chaperone that inhibits aggregation of misfolded proteins. At low concentrations, facilitates aggregation (anti-chaperone activity). May be involved with other chaperones in the structural modification of the TG precursor in hormone biogenesis. Also acts a structural subunit of various enzymes such as prolyl 4-hydroxylase and microsomal triacylglycerol transfer protein MTTP. Receptor for LGALS9; the interaction retains P4HB at the cell surface of Th2 T helper cells, increasing disulfide reductase activity at the plasma membrane, altering the plasma membrane redox state and enhancing cell migration (PubMed:21670307).

Synonyms:

DSI; ERBA2L; GIT; P4Hbeta; p55; PDI; PDIA1; PHDB; PO4DB; PO4HB; PROHB

Product images:


Western blot analysis of P4HB in K562, C6 lysates using P4HB antibody.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using P4HB antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.