

## Product datasheet for **TA384918S**

### **P4HB Rabbit Monoclonal Antibody [Clone ID: R08-6I8]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	R08-6I8
Applications:	WB
Recommended Dilution:	WB: 1/2000-1/10000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Immunogen:	A synthetic peptide of human PDIA2
Formulation:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year
Predicted Protein Size:	Calculated MW: 58 kDa; Observed MW: 58 kDa
Gene Name:	prolyl 4-hydroxylase subunit beta
Database Link:	<a href="#">Entrez Gene 5034 Human P07237</a>




[View online »](#)

**Background:**

Swiss-Prot Acc.P07237. This multifunctional protein catalyzes the formation, breakage and rearrangement of disulfide bonds. At the cell surface, seems to act as a reductase that cleaves disulfide bonds of proteins attached to the cell. May therefore cause structural modifications of exofacial proteins. Inside the cell, seems to form/rearrange disulfide bonds of nascent proteins. At high concentrations, functions as a chaperone that inhibits aggregation of misfolded proteins. At low concentrations, facilitates aggregation (anti-chaperone activity). May be involved with other chaperones in the structural modification of the TG precursor in hormone biogenesis. Also acts a structural subunit of various enzymes such as prolyl 4-hydroxylase and microsomal triacylglycerol transfer protein MTTP. Receptor for LGALS9; the interaction retains P4HB at the cell surface of Th2 T helper cells, increasing disulfide reductase activity at the plasma membrane, altering the plasma membrane redox state and enhancing cell migration (PubMed:21670307).

**Synonyms:**

DSI; ERBA2L; GIT; P4Hbeta; p55; PDI; PDIA1; PHDB; PO4DB; PO4HB; PROHB

**Product images:**

Western blot analysis of PDIA2 in K562, C6 lysates using P4HB antibody.