

Product datasheet for TA384925S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PAK2 Mouse Monoclonal Antibody [Clone ID: 7H3-E2-F5]

Product data:

Product Type: Primary Antibodies

Clone Name: 7H3-E2-F5
Applications: IF, IHC, WB

Recommended Dilution: WB: 1/500-1/2000

IHC: 1/200-1/1000 ICC: 1/200-1/1000

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Purified recombinant fragment of PAK2 expressed in E. Coli.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.03% Proclin 300, pH 7.3.

Concentration:lot specificPurification:Ascitic FluidConjugation:Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year Predicted Protein Size: 61kDa

Gene Name: p21 (RAC1) activated kinase 2

Database Link: Entrez Gene 5062 Human

Q13177

Background: Swiss-Prot Acc.Q13177.PAK2, also known as P21 (CDKN1A)-activated kinase 2. The p21

activated kinases (PAK) are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. The PAK proteins are a family of serine/threonine kinases that serve as targets for the small GTP binding proteins, CDC42 and RAC1, and have been implicated in a wide range of biological activities. PAK2 is activated by proteolytic cleavage during caspase-mediated apoptosis, and may play a role in regulating the apoptotic events in the dying cell. PAK2 has been shown to interact with SH3KBP1, CDC42 and Abl gene.




Synonyms:

Gamma-PAK; p58; PAK-2; PAK65; PAKgamma


Product images:



Confocal immunofluorescence analysis of Hela cells using PAK2 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye.



Western blot detection of PAK2 in Jurkat,MCF7 and Hela cell lysates using PAK2 mouse mAb (1:5000 diluted).Predicted band size:61KDa.Observed band size:61KDa.



Immunohistochemical analysis of paraffinembedded human lung cancer (left) and gastric cancer (right) using PAK2 mouse mAb with DAB staining.