

Product datasheet for TA384981M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PI 3 Kinase p85 alpha (PIK3R1) Mouse Monoclonal Antibody [Clone ID: 1C8-2H4-9D9]

Product data:

Product Type: Primary Antibodies

Clone Name: 1C8-2H4-9D9

Applications: IHC, WB

Recommended Dilution: WB: 1/1000-2000

IHC: 1/100-200

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Purified recombinant protein expressed in E.coli.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.03% Proclin 300, pH 7.3.

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year
Predicted Protein Size: 85kDa

Gene Name: phosphoinositide-3-kinase regulatory subunit 1

Database Link: Entrez Gene 5295 Human

P27986






Background:


Swiss-Prot Acc.P27986.Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:17626883, PubMed:19805105, PubMed:7518429). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed:20348923).

Synonyms: GRB1; p85; p85-ALPHA; PI3K

Product images:




Western blot detection of PI3 Kinase p85 α (1C8) in Hela,A549,HL-60,U2OS,C6 cell lysates using PI3 Kinase p85 α (1C8) antibody. Observed band size:85kDa.




Western blot analysis of PI3 Kinase p85 alpha (1C8) in Hela, A549, HL-60, U2OS, C6 lysates using PI3 Kinase p85 alpha (1C8) antibody.





Immunohistochemistry analysis of paraffinembedded rat Liver Tissue using PI3 Kinase p85 alpha (1C8) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffinembedded Human tonsils using PI3 Kinase p85 alpha (1C8) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.