

# **Product datasheet for TA384989M**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## PKN1 Rabbit Monoclonal Antibody [Clone ID: R09-214]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: R09-214

Applications: WB

Recommended Dilution: WB: 1/1000-1/5000

**Reactivity:** Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

**Immunogen:** A synthetic peptide of human PKN

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 104 kDa; Observed MW: 120 kDa

**Gene Name:** protein kinase N1

Database Link: Q16512




#### Background:

Swiss-Prot Acc.Q16512.PKC-related serine/threonine-protein kinase involved in various processes such as regulation of the intermediate filaments of the actin cytoskeleton, cell migration, tumor cell invasion and transcription regulation. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14. Regulates the cytoskeletal network by phosphorylating proteins such as VIM and neurofilament proteins NEFH, NEFL and NEFM, leading to inhibit their polymerization. Phosphorylates 'Ser-575', 'Ser-637' and 'Ser-669' of MAPT/Tau, lowering its ability to bind to microtubules, resulting in disruption of tubulin assembly. Acts as a key coactivator of androgen receptor (ANDR)-dependent transcription, by being recruited to ANDR target genes and specifically mediating phosphorylation of 'Thr-11' of histone H3 (H3T11ph), a specific tag for epigenetic transcriptional activation that promotes demethylation of histone H3 'Lys-9' (H3K9me) by KDM4C/JMJD2C. Phosphorylates HDAC5, HDAC7 and HDAC9, leading to impair their import in the nucleus. Phosphorylates 'Thr-38' of PPP1R14A, 'Ser-159', 'Ser-163' and 'Ser-170' of MARCKS, and GFAP. Able to phosphorylate RPS6 in vitro.

Synonyms:

DBK; MGC46204; PAK1; PKN; PKN-ALPHA; PRK1; PRKCL1

### **Product images:**



Western blot analysis of PKN in C6 lysates using PKN1 antibody.