

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA385050

### PKC beta 1 (PRKCB) Rabbit Monoclonal Antibody [Clone ID: R06-0C2]

#### **Product data:**

| Product Type:           | Primary Antibodies   |
|-------------------------|--|
| Clone Name:             | R06-0C2  |
| Applications:           | IF, IHC, IP, WB  |
| Recommended Dilution:   | WB: 1/500-1/2000<br>IHC: 1/50-1/200<br>ICC/IF: 1/50-1/200<br>IP: 1/50                    |
| Reactivity:             | Human, Rat   |
| Host:                   | Rabbit   |
| lsotype:                | IgG  |
| Clonality:              | Monoclonal   |
| Immunogen:              | A synthesized peptide derived from human PKC beta 2                                      |
| Formulation:            | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA    |
| Concentration:          | lot specific   |
| Purification:           | Affinity Purified  |
| Conjugation:            | Unconjugated   |
| Storage:                | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Stability:              | 1 year   |
| Predicted Protein Size: | 77kDa  |
| Gene Name:              | protein kinase C beta  |
| Database Link:          | <u>Entrez Gene 5579 Human</u><br><u>P05771</u>   |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

|             | PKC beta 1 (PRKCB) Rabbit Monoclonal Antibody [Clone ID: R06-0C2] – TA385050  |
|-------------|---|
| Background: | Swiss-Prot Acc.P05771.Calcium-activated, phospholipid- and diacylglycerol (DAG)-dependent<br>serine/threonine-protein kinase involved in various cellular processes such as regulation of<br>the B-cell receptor (BCR) signalosome, oxidative stress-induced apoptosis, androgen<br>receptor-dependent transcription regulation, insulin signaling and endothelial cells<br>proliferation. Plays a key role in B-cell activation by regulating BCR-induced NF-kappa-B<br>activation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by direct<br>phosphorylation of CARD11/CARMA1 at 'Ser-559', 'Ser-644' and 'Ser-652'. |

#### Synonyms: MGC41878; PKC-B; PKC-beta; PKCB; PRKCB1; PRKCB2


## **Product images:**




Western blot analysis of PKC beta 2 in rat brain lysates using PKC beta 2 antibody.Observed band size:77kDa.


Western blot analysis of PKC beta 2 in Hela, A549, U2OS, U87-MG lysates using PKC beta 2 antibody

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry analysis of paraffinembedded Human tonsil using PKC beta 2 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunocytochemistry analysis of PKC beta 2 (green) in K562 using PKC beta 2 antibody,and DAPI(blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US