

# **Product datasheet for TA385067M**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

### PSMB8 Mouse Monoclonal Antibody [Clone ID: 3G3-C1-E2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: 3G3-C1-E2
Applications: IF, IHC, WB

Recommended Dilution: WB: 1/500-1/2000

IHC: 1/200-1/1000 ICC: 1/200-1/1000

Reactivity: Human, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Purified recombinant fragment of human PSMB8 expressed in E. Coli.

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.03% Proclin 300, pH 7.3.

Concentration: lot specific

Purification: Ascitic Fluid

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year Predicted Protein Size: 30kDa

Gene Name: proteasome subunit beta 8

Database Link: Entrez Gene 5696 Human

P28062






#### Background:

Swiss-Prot Acc.P28062. The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to yield the same mature subunit.

Synonyms:


beta5i; D6S216; D6S216E; LMP7; MGC1491; PSMB5i; RING10; Y2

## **Product images:**




Immunofluorescence analysis of Hela cells using PSMB8 mouse mAb (green). Blue





Western blot analysis of PSMB8/LMP7 in C6, Hela, Molt4 and Raji lysates using PSMB8/LMP7 antibody.



Immunohistochemical analysis of paraffinembedded intima cancer tissues using PSMB8 mouse mAb with DAB staining.