

Product datasheet for TA385140

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rb (RB1) Rabbit Monoclonal Antibody [Clone ID: R01-8H7]

Product data:

Product Type: Primary Antibodies

Clone Name: R01-8H7
Applications: IP, WB

Recommended Dilution: WB: 1/1000

IP: 1/20

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic phosphopeptide corresponding to residues surrounding Ser780 of human Rb

(Phosphorylated)

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 106 kDa; Observed MW: 110 kDa

Gene Name: RB transcriptional corepressor 1

Database Link: Entrez Gene 5925 Human

P06400






Background:

Swiss-Prot Acc.P06400.Key regulator of entry into cell division that acts as a tumor suppressor. Promotes G0-G1 transition when phosphorylated by CDK3/cyclin-C. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. Mediates transcriptional repression by SMARCA4/BRG1 by recruiting a histone deacetylase (HDAC) complex to the c-FOS promoter. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC1 repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex .

Synonyms:

OSRC; osteosarcoma; p105-Rb; pp110; pRb; RB

Product images:



Western blot analysis of Phospho-Rb (Ser780) in K562, C6 lysates using Phospho-Rb (Ser780) antibody.