

Product datasheet for TA385146

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

hnRNP G (RBMX) Rabbit Monoclonal Antibody [Clone ID: R03-1C9]

Product data:

Product Type: Primary Antibodies

Clone Name: R03-1C9
Applications: IF, IHC, WB
Recommended Dilution: WB: 1/1000

IHC: 1/20-1/100 ICC/IF: 1/20-1/50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human hnRNP G

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 42 kDa; Observed MW: 42 kDa

Gene Name: RNA binding motif protein, X-linked

Database Link: Entrez Gene 27316 Human

P38159






Background:

Swiss-Prot Acc.P38159.RNA-binding protein that plays several role in the regulation of preand post-transcriptional processes. Implicated in tissue-specific regulation of gene transcription and alternative splicing of several pre-mRNAs. Binds to and stimulates transcription from the tumor suppressor TXNIP gene promoter; may thus be involved in tumor suppression. When associated with SAFB, binds to and stimulates transcription from the SREBF1 promoter. Associates with nascent mRNAs transcribed by RNA polymerase II. Component of the supraspliceosome complex that regulates pre-mRNA alternative splice site selection. Can either activate or suppress exon inclusion; acts additively with TRA2B to promote exon 7 inclusion of the survival motor neuron SMN2. Represses the splicing of MAPT/Tau exon 10. Binds preferentially to single-stranded 5'-CC[A/C]-rich RNA sequence motifs localized in a single-stranded conformation; probably binds RNA as a homodimer. Binds non-specifically to pre-mRNAs. Plays also a role in the cytoplasmic TNFR1 trafficking pathways; promotes both the IL-1-beta-mediated inducible proteolytic cleavage of TNFR1 ectodomains and the release of TNFR1 exosome-like vesicles to the extracellular compartment.

Synonyms:


hnRNP-G; HNRPG; RBMXP1; RBMXRT; RNMX

Product images:




Western blot analysis of hnRNP G in K562, Hela lysates using hnRNP G antibody.





Immunohistochemistry analysis of paraffinembedded Human lung cancer tissue using hnRNP G antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunocytochemistry analysis of hnRNP G (green) in 293T using hnRNP G antibody,and DAPI(blue).