

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA385148

RBPJK (RBPJ) Rabbit Monoclonal Antibody [Clone ID: R05-6G4]

Product data:

| Clone Name:R05-6G4Applications:WBRecommended Dilution:WB: 1/1000-1/5000Reactivity:Rat |
|--|
| Recommended Dilution: WB: 1/1000-1/5000 |
| |
| Reactivity: Rat |
| |
| Host: Rabbit |
| lsotype: lgG |
| Clonality: Monoclonal |
| Immunogen: A synthetic peptide of human RBPJK |
| Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Concentration: lot specific |
| Purification: Affinity Purified |
| Conjugation: Unconjugated |
| Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Stability: 1 year |
| Predicted Protein Size: Calculated MW: 56 kDa; Observed MW: 61 kDa |
| Gene Name: recombination signal binding protein for immunoglobulin kappa J region |
| Database Link: <u>Q06330</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


SRIGENE RBPJK (RBPJ) Rabbit Monoclonal Antibody [Clone ID: R05-6G4] – TA385148

Background: Swiss-Prot Acc.Q06330.Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins (Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA (PubMed:21991380). Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen) (PubMed:23303788). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase subunits .

 Synonyms:
 CBF-1; CBF1; csl; IGKJRB; IGKJRB1; KBF2; MGC61669; OTTHUMP00000123441; RBP-J; RBP-JK;

 RBPJK; RBPSUH; SUH

Product images:



Western blot analysis of RBPJK in rat Brain, C6 lysates using RBPJK antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US