

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA385148S

RBPJK (RBPJ) Rabbit Monoclonal Antibody [Clone ID: R05-6G4]

Product data:

Clone Name: R05-6G4	
Applications: WB	
Recommended Dilution: WB: 1/1000-1/5000	
Reactivity: Rat	
Host: Rabbit	
lsotype: lgG	
Clonality: Monoclonal	
Immunogen: A synthetic peptide of human RBPJK	
Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium a	zide and 0.05% BSA
Concentration: lot specific	
Purification: Affinity Purified	
Conjugation: Unconjugated	
Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid free	eeze/thaw cycles.
Stability: 1 year	
Predicted Protein Size: Calculated MW: 56 kDa; Observed MW: 61 kDa	
Gene Name: recombination signal binding protein for immunoglobulin kappa J regi	on
Database Link: <u>Q06330</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Scherker Content of the second state of the se

Background: Swiss-Prot Acc.Q06330.Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins (Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA (PubMed:21991380). Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen) (PubMed:23303788). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase subunits .

 Synonyms:
 CBF-1; CBF1; csl; IGKJRB; IGKJRB1; KBF2; MGC61669; OTTHUMP00000123441; RBP-J; RBP-JK;

 RBPJK; RBPSUH; SUH

Product images:



Western blot analysis of RBPJK in rat Brain, C6 lysates using RBPJK antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US