

# **Product datasheet for TA385205**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### S100P Mouse Monoclonal Antibody [Clone ID: 7C5-6G8-5F3]

#### **Product data:**

**Product Type:** Primary Antibodies

**Clone Name:** 7C5-6G8-5F3

Applications: IHC

Recommended Dilution: IHC-p: 1/100-1/500

Reactivity: Human Host: Mouse

Isotype: IgG2a, kappa
Clonality: Monoclonal

Immunogen: Synthesized peptide derived from human S100P

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

**Concentration:** lot specific

**Purification:** Affinity Chromatography

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

**Gene Name:** S100 calcium binding protein P

**Database Link:** Entrez Gene 6286 Human


P25815

**Background:** Swiss-Prot Acc.P25815.


Synonyms: MIG9; S100E



## **Product images:**



Immunohistochemical analysis of paraffinembedded Pancreatic carcinoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffinembedded Pancreatic carcinoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





Immunohistochemical analysis of paraffinembedded Pancreatic carcinoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, TRIS-EDTA of pH8. was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffinembedded Pancreatic carcinoma. 1, Antibody was diluted at 1:200(4°,overnight). 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).