

Product datasheet for TA385338M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STAT3 Rabbit Monoclonal Antibody [Clone ID: R05-7B2]

Product data:

Product Type: Primary Antibodies

Clone Name: R05-7B2
Applications: IF, WB

Recommended Dilution: WB: 1/1000

ICC/IF: 1/20

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic phosphopeptide corresponding to residues surrounding Ser727 of human STAT3

(Phosphorylated)

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 88 kDa; Observed MW: 88 kDa

Gene Name: signal transducer and activator of transcription 3

Database Link: Entrez Gene 6774 Human

P40763






Background:

Swiss-Prot Acc.P40763.Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors (PubMed:10688651, PubMed:12359225, PubMed:12873986, PubMed:15194700, PubMed:17344214, PubMed:18242580, PubMed:23084476). Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed:17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed:12873986). Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes (PubMed:12359225). Activated by IL31 through IL31RA (PubMed:15194700). Acts as a regulator of inflammatory response by regulating differentiation of naive CD4+ Tcells into T-helper Th17 or regulatory T-cells (Treg): deacetylation and oxidation of lysine residues by LOXL3, leads to disrupt STAT3 dimerization and inhibit its transcription activity (PubMed:28065600). Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation. May play an apoptotic role by transctivating BIRC5 expression under LEP activation (PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity (PubMed:23084476). Plays a crucial role in basal beta cell functions, such as regulation of insulin secretion.

Synonyms: APRF; FLJ20882; HIES; MGC16063

Product images:



Western blot analysis of Phospho-STAT3 (Ser727) in Jurkat, C6, Hela lysates using Phospho-STAT3 (Ser727) antibody.