

# **Product datasheet for TA385363**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **SUN1 Rabbit Monoclonal Antibody [Clone ID: R05-9D5]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R05-9D5

Applications: IF, IHC, WB

Recommended Dilution: WB: 1/1000

IHC: 1/50 ICC/IF: 1/50

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

**Immunogen:** A synthetic peptide of human SUN1

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 90 kDa; Observed MW: 90 kDa

Gene Name: Sad1 and UNC84 domain containing 1

Database Link: Entrez Gene 23353 Human

O94901






#### Background:

Swiss-Prot Acc. O94901. As a component of the LINC (LInker of Nucleoskeleton and Cytoskeleton) complex involved in the connection between the nuclear lamina and the cytoskeleton (PubMed:18039933, PubMed:18396275). The nucleocytoplasmic interactions established by the LINC complex play an important role in the transmission of mechanical forces across the nuclear envelope and in nuclear movement and positioning. Required for interkinetic nuclear migration (INM) and essential for nucleokinesis and centrosome-nucleus coupling during radial neuronal migration in the cerebral cortex and during glial migration. Involved in telomere attachment to nuclear envelope in the prophase of meiosis implicating a SUN1/2:KASH5 LINC complex in which SUN1 and SUN2 seem to act at least partial redundantly. Required for gametogenesis and involved in selective gene expression of coding and non-coding RNAs needed for gametogenesis. Helps to define the distribution of nuclear pore complexes (NPCs). Required for efficient localization of SYNE4 in the nuclear envelope. May be involved in nuclear remodeling during sperm head formation in spermatogenenis. May play a role in DNA repair by suppressing non-homologous end joining repair to facilitate the repair of DNA cross-links (PubMed:24375709).


Synonyms: UNC84A

## **Product images:**




Western blot analysis of SUN1 in K562, Hela lysates using SUN1 antibody.





Immunohistochemistry analysis of paraffinembedded Human breast cancer using SUN1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunocytochemistry analysis of SUN1(green) in Hela using SUN1 antibody, and DAPI(blue)