

## **Product datasheet for TA385408S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TRIF (TICAM1) Rabbit Monoclonal Antibody [Clone ID: R02-2A9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R02-2A9
Applications: IP, WB

Recommended Dilution: WB: 1/1000

IP: 1/20

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: Recombinant protein of human TRIF

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

**Storage:** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 76 kDa; Observed MW: 98 kDa

**Gene Name:** toll like receptor adaptor molecule 1

Database Link: Entrez Gene 148022 Human

Q8IUC6




#### Background:


Swiss-Prot Acc.Q8IUC6.Involved in innate immunity against invading pathogens. Adapter used by TLR3 and TLR4 (through TICAM2) to mediate NF-kappa-B and interferon-regulatory factor (IRF) activation, and to induce apoptosis. Ligand binding to these receptors results in TRIF recruitment through its TIR domain. Distinct protein-interaction motifs allow recruitment of the effector proteins TBK1, TRAF6 and RIPK1, which in turn, lead to the activation of transcription factors IRF3 and IRF7, NF-kappa-B and FADD respectively. Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of proinflammatory cytokines .

Synonyms: MGC35334; PRVTIRB; TICAM-1; TRIF

# **Product images:**



Western blot analysis of TRIF in rat testis lysates using TRIF antibody.



Western blot analysis of TRIF in K562, 3T3, Hela lysates using TRIF antibody.