

Product datasheet for TA385447M

Trp53 Rabbit Monoclonal Antibody [Clone ID: R01-2B9]

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Clone Name:	R01-2B9
Applications:	IF, IP, WB
Recommended Dilution:	WB: 1/1000 ICC/IF: 1/20 IP: 1/20
Reactivity:	Mouse
Host:	Rabbit
lsotype:	IgG
Clonality:	Monoclonal
Immunogen:	Recombinant protein of mouse p53
Formulation:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year
Predicted Protein Size:	Calculated MW: 44 kDa; Observed MW: 53 kDa
Gene Name:	transformation related protein 53
Database Link:	<u>Entrez Gene 22059 Mouse</u> <u>P02340</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE Trp53 Rabbit Monoclonal Antibody [Clone ID: R01-2B9] – TA385447M

Background:Swiss-Prot Acc.P02340.Acts as a tumor suppressor in many tumor types; induces growth
arrest or apoptosis depending on the physiological circumstances and cell type. Involved in
cell cycle regulation as a trans-activator that acts to negatively regulate cell division by
controlling a set of genes required for this process. One of the activated genes is an inhibitor
of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation
of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with
mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is
largely independent of transcription. Prevents CDK7 kinase activity when associated to CAK
complex in response to DNA damage, thus stopping cell cycle progression . Induces the
transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1.
LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis,
but seems to have to effect on cell-cycle regulation. Regulates the circadian clock by
repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2
(PubMed:24051492).

Synonyms:

bbl; bfy; bhy; p44; p53; Tp53

Product images:



Western blot analysis of p53 in 3T3 lysates using p53 antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US