

Product datasheet for TA385539M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VCAM1 Rabbit Monoclonal Antibody [Clone ID: R02-7H1]

Product data:

Product Type: Primary Antibodies

Clone Name: R02-7H1

Applications: IHC, IP, WB

Recommended Dilution: WB: 1/2000

IHC: 1/20-1/500

IP: 1/20

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of mouse VCAM1

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 81 kDa; Observed MW: 110 kDa

Gene Name: vascular cell adhesion molecule 1

Database Link: Entrez Gene 7412 Human

P19320

Background: Swiss-Prot Acc.P19320.Important in cell-cell recognition. Appears to function in leukocyte-

endothelial cell adhesion. Interacts with integrin alpha-4/beta-1 (ITGA4/ITGB1) on leukocytes, and mediates both adhesion and signal transduction. The VCAM1/ITGA4/ITGB1 interaction may play a pathophysiologic role both in immune responses and in leukocyte emigration to


sites of inflammation.

Synonyms: CD106; DKFZp779G2333; INCAM-100; L1CAM; MGC99561






Product images:




Western blot analysis of VCAM1 in rat spleen, rat kidney lysates using VCAM1 antibody.



Western blot analysis of VCAM1 in 3T3 lysates using VCAM1 antibody.





Immunohistochemistry analysis of paraffinembedded Human tonsil using VCAM1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.