

Product datasheet for TA385554

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VPS4A Rabbit Monoclonal Antibody [Clone ID: R01-9H9]

Product data:

Product Type: Primary Antibodies

Clone Name: R01-9H9
Applications: IF, IHC, WB

Recommended Dilution: WB: 1/2000-1/10000

IHC: 1/500 ICC/IF: 1/100

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human VPS4a

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 49 kDa; Observed MW: 49 kDa

Gene Name: vacuolar protein sorting 4 homolog A

Database Link: Entrez Gene 27183 Human

Q9UN37




Background:

Swiss-Prot Acc.Q9UN37.Involved in late steps of the endosomal multivesicular bodies (MVB) pathway. Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly, possibly in combination with membrane fission. Redistributes the ESCRT-III components to the cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. In conjunction with the ESCRT machinery also appears to function in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and enveloped virus budding (HIV-1 and other lentiviruses). Involved in cytokinesis: retained at the midbody by ZFYVE19/ANCHR and CHMP4C until abscission checkpoint signaling is terminated at late cytokinesis. It is then released following dephosphorylation of CHMP4C, leading to abscission (PubMed:24814515). VPS4A/B are required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:22660413).

Synonyms:


FLJ22197; hVPS4; SKD1; SKD1-homolog; SKD1A; SKD2; VPS4; VPS4-1

Product images:




Western blot analysis of VPS4a in K562, rat Brain, C6, 3T3 lysates using VPS4A antibody.





Immunohistochemistry analysis of paraffinembedded Human breast cancer using VPS4a antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunocytochemistry analysis of VPS4a(green) in Hela using VPS4a antibody,and DAPI(blue)