

Product datasheet for TA385570M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850 LIS

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

YAP1 Rabbit Monoclonal Antibody [Clone ID: R06-6H7]

Product data:

Product Type: Primary Antibodies

Clone Name: R06-6H7
Applications: IHC, IP, WB

Recommended Dilution: WB: 1/2000-1/10000

IHC: 1/20-1/100

IP: 1/20

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human YAP1

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 55 kDa; Observed MW: 65 kDa

Gene Name: Yes associated protein 1

Database Link: Entrez Gene 10413 Human

P46937






Background:

Swiss-Prot Acc.P46937.Transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis (PubMed:17974916, PubMed:18280240, PubMed:18579750, PubMed:21364637). The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ (PubMed:18158288). Plays a key role in tissue tension and 3D tissue shape by regulating cortical actomyosin network formation. Acts via ARHGAP18, a Rho GTPase activating protein that suppresses Factin polymerization (PubMed:25778702). Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration (PubMed:18158288). The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction (PubMed:18579750).


Synonyms: YAP; YAP2; YAP65; YKI

Product images:



Western blot analysis of YAP1 in Hela lysates using YAP1 antibody.





Immunohistochemistry analysis of paraffinembedded Human lung cancer using YAP1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.