OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for TA385598

Actin Mouse Monoclonal Antibody [Clone ID: 8B10-3F7-1B8]

Product data:

Product Type:	Primary Antibodies
Clone Name:	8B10-3F7-1B8
Applications:	IHC
Recommended Dilution:	IHC-p: 1/100-1/500
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1, kappa
Clonality:	Monoclonal
Immunogen:	Synthesized peptide derived from human Actin, sarcomeric muscle
Formulation:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Concentration:	lot specific
Purification:	Affinity Chromatography
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year

View online »

Product images:



Immunohistochemical analysis of paraffin-embedded human Myocardium. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval



Immunohistochemical analysis of paraffin-embedded human Myocardium. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval



Immunohistochemical analysis of paraffin-embedded human Myocardium. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6. was used for antigen retrieval



Immunohistochemical analysis of paraffin-embedded human Prostate. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval



Immunohistochemical analysis of paraffin-embedded human Skeletal muscle. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval



Immunohistochemical analysis of paraffin-embedded human Smooth muscle. 1,Actin, sarcomeric muscle Antibody was diluted at 1:200(4°,overnight). 2, Citrate buffer of pH6.0 was used for antigen retrieval