

Product datasheet for TA385892

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HSP27 (HSPB1) Rabbit Monoclonal Antibody [Clone ID: SAIC-23C-35]

Product data:

Product Type: Primary Antibodies

Clone Name: SAIC-23C-35

Applications: WB

Reactivity: Human Host: Rabbit

Isotype: IgG, kappa
Clonality: Monoclonal

Immunogen: Peptide "VSLDVNHFAPDELTVK" derived from Heat shock protein beta-1, involved in stress

resistance and actin organisation, conjugated to KLH.

Specificity: Recognises human HSPB1.

Original data characterizing this antibody may be foundhere.

Formulation: PBS with 0.02% Proclin 300.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Please store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C. Avoid

freeze and thaw cycles.

Stability: 3 years from dispatch.

Gene Name: heat shock protein family B (small) member 1


Database Link: Entrez Gene 3315 Human

P04792


Synonyms: CMT2F; DKFZp586P1322; HMN2B; HS.76067; Hsp25; HSP27; HSP28; SRP27



Product images:



Flow cytometry using the Anti-HSPB1 antibody SAIC-23C-35 (TA385892). Paraformaldehyde fixed HeLa cells permeabilized with 0.5% Triton were stained with anti-unknown specificity antibody ([TA385792]; isotype control, black line) or the rabbit IgG version of SAIC-23C-35 (TA385892, blue line) at a dilution of 1:100 for 1h at RT. After washing, the bound antibody was detected using a goat anti-rabbit IgG AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.



Western Blot using anti-HSPB1 antibody SAIC-23C-35 (TA385892). HeLa(A) (0.03µg/ml), MCF7(B) (0.001µg/ml), Caco-2(C) (0.1µg/ml) and A431(D) (0.03µg/ml) cell lysates (35µg protein in RIPA buffer) were resolved on a SDS PAGE gel and blots were probed with the chimeric rabbit version of SAIC-23C-35 (TA385892), before detection using an anti-rabbit secondary antibody. A primary incubation of 1h was used and protein was detected by chemiluminescence.