

Product datasheet for TA386536

CD28 Mouse Monoclonal Antibody [Clone ID: 150000000]

Primary Antibodies

150000000

ELISA, FC

Human

Mouse

IgG1, kappa

Monoclonal

The original mouse antibody was generated by immunizing mice with human T lymphocytes.

Product data:

Product Type:

Clone Name:

Applications:

Reactivity:

Host:

Isotype:

Clonality:

Immunogen:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CD28 Mouse Monoclonal Antibody [Clone ID: 1500000000] – TA386536
Specificity:	This antibody binds human CD28, which is found on a sub-population of T cells and activated B cells. The antibody does not react with B cells, granulocytes or monocytes. CD28 is involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival.
	This antibody works independently of TcR/CD3 and activation of protein kinase C (PKC) for activation of T cells and nehances cytokine secretion (PMID: 1647959). This antibody is capable of stimulating T cells in combination with either murine OKT3 antibody or CDR-mutated OKT3 Fab variants. It was found that cross-linking of specific tumor antigens with the T-cell-associated CD3 and CD28 antigens can increase IL-2 secretion, proliferation and antigen-specific cytotxicity in resting T cells. This cross-linking can be achieved effectively by bispecific monoclonal antibodies (BiMAb) with specificity for both the tumor antigen and CD3 or CD28 antigen. A combination of bispecific antibodies BiMAb OKT3/HRS-3 with reactivity against both CD3 and Hodgkin's-lymphoma-associated CD30 antigen and BiMAb 15E8/HRS-3 with reactivity to both CD28 and CD30 antigen was able to activate resting T cells, represented by Jurkat cells (CD3+/CD28+) to produce IL-2 (PMID: 7686889). This antibody was also used in another study wherein the combined use of two CD3 x CD19 plus CD28 x CD22 bispecific antibodies induced optimal interleukin 2 secretion by Jurkat T-cell acute lymphocytic leukemia cells in the presence of target B-cells (PMID: 7689932). This antibody also provides a co-stimulatory signal to T cells. The scFv fragment of this antibody induced proliferation of peripheral blood mononuclear cells in the presence of anti-CD3 or anti-CD2 mAb and enhanced anti-tumor activity of anti-MUC1×anti-CD3 bispecific antibody when tested with lymphokine-activated killer cells with T cell phenotype (PMID: 10913626). The humanzied version of this antibody is available on request.
Formulation:	PBS with 0.02% Proclin 300.
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Please store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C. Avoid freeze and thaw cycles.
Stability:	3 years from dispatch.
Gene Name:	CD28 molecule
Database Link:	<u>Entrez Gene 940 Human</u> <u>P10747</u>
Synonyms:	MGC138290; Tp44

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US