

## **Product datasheet for TA386644**

**H1-2 Rabbit Polyclonal Antibody** 

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:100-1:1000, IHC:1:20-1:200

**Reactivity:** Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Peptide sequence around site of Lys (33) derived from Human Histone H1.2

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: P16403

**Background:** Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular

structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation

(By similarity).




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded human lung cancer using TA386644 at dilution of 1:100



Western Blot

Positive WB detected in: MCF-7 whole cell lysate, Mouse brain tissue, Mouse liver tissue All lanes: HIST1H1C antibody at 0.3µg/ml Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 22 kDa

Observed band size: 22 kDa