

Product datasheet for **TA386654**


H1-2 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	ICC, IF, WB
Recommended Dilution:	Recommended dilution: WB:1:100-1:1000, ICC:1:10-1:100, IF:1:10-1:100
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide sequence around site of Formyl-Lys (109) derived from Human Histone H1.2
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Concentration:	lot specific
Purification:	Antigen Affinity Purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	P16403
Background:	Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).




[View online »](#)

Product images:

Immunofluorescence staining of HeLa cells with TA386654 at 1:20, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunocytochemistry analysis of TA386654 diluted at 1:20 and staining in HeLa cells (treated with 30mM sodium butyrate for 4h) performed on a Leica Bond™ system. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot
 Detected samples: K562 whole cell lysate;
 Untreated (-) or treated (+) with 30mM sodium
 butyrate for 4h
 All lanes: HIST1H1C antibody at 1:400
 Secondary
 Goat polyclonal to rabbit IgG at 1/50000 dilution
 Predicted band size: 22 kDa
 Observed band size: 22 kDa