

Product datasheet for TA386693M

Product data:

Product Type: Primary Antibodies

USP8 Rabbit Polyclonal Antibody

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Ubiquitin carboxyl-terminal hydrolase 8 protein (1-270AA)

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Antigen Affinity Purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: P40818

Background: Hydrolase that can remove conjugated ubiquitin from proteins and therefore plays an

important regulatory role at the level of protein turnover by preventing degradation.

Converts both 'Lys-48' an 'Lys-63'-linked ubiquitin chains. Catalytic activity is enhanced in the M phase. Involved in cell proliferation. Required to enter into S phase in response to serum stimulation. May regulate T-cell anergy mediated by RNF128 via the formation of a complex containing RNF128 and OTUB1. Probably regulates the stability of STAM2 and RASGRF1. Regulates endosomal ubiquitin dynamics, cargo sorting, membrane traffic at early endosomes, and maintenance of ESCRT-0 stability. The level of protein ubiquitination on endosomes is essential for maintaining the morphology of the organelle. Deubiquitinates

EPS15 and controles tyrosine kinase stability. Removes conjugated ubiquitin from EGFR thus

regulating EGFR degradation and downstream MAPK signaling. Involved in acrosome biogenesis through interaction with the spermatid ESCRT-0 complex and microtubules.

Deubiquitinates BIRC6/bruce and KIF23/MKLP1.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Immunohistochemistry of paraffin-embedded human testis tissue using [TA386693] at dilution of 1:100



Western blot

All lanes: USP8 antibody at $14\mu g/ml + 293T$ whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 128, 116 kDa Observed band size: 128 kDa