

## **Product datasheet for TA386695M**

## Troduct datasficct for TASSOUSSIN

## **GRK6 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200

**Reactivity:** Rat, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human G protein-coupled receptor kinase 6 protein (1-170AA)

**Formulation:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified

Conjugation: Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: P43250

**Background:** Specifically phosphorylates the activated forms of G protein-coupled receptors. Such receptor

phosphorylation initiates beta-arrestin-mediated receptor desensitization, internalization,

and signaling events leading to their desensitization. Seems to be involved in the

desensitization of D2-like dopamine receptors in striatum and chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis (By similarity). Phosphorylates rhodopsin (RHO) (in vitro) and a non G-protein-coupled receptor: LRP6 during Wnt signaling (in vitro).




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded human kidney tissue using [TA386695] at dilution of 1:100



Western blot
All lanes: G protein-coupled receptor kinase 6
antibody at 6µg/ml + Rat heart tissue
Secondary
Goat polyclonal to rabbit IgG at 1/10000 dilution
Predicted band size: 66, 68, 65 kDa
Observed band size: 66 kDa