

Product datasheet for TA386860M

ITPK1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Inositol-tetrakisphosphate 1-kinase protein (185-414AA)

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Antigen Affinity Purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q13572

Background: Kinase that can phosphorylate various inositol polyphosphate such as Ins(3,4,5,6)P4 or

Ins(1,3,4)P3. Phosphorylates Ins(3,4,5,6)P4 at position 1 to form Ins(1,3,4,5,6)P5. This reaction is thought to have regulatory importance, since Ins(3,4,5,6)P4 is an inhibitor of plasma membrane Ca(2+)-activated Cl(-) channels, while Ins(1,3,4,5,6)P5 is not. Also phosphorylates

Ins(1,3,4)P3 on O-5 and O-6 to form Ins(1,3,4,6)P4, an essential molecule in the

hexakisphosphate (InsP6) pathway. Also acts as an inositol polyphosphate phosphatase that dephosphorylate Ins(1,3,4,5)P4 and Ins(1,3,4,6)P4 to Ins(1,3,4)P3, and Ins(1,3,4,5,6)P5 to Ins(3,4,5,6)P4. May also act as an isomerase that interconverts the inositol tetrakisphosphate isomers Ins(1,3,4,5)P4 and Ins(1,3,4,6)P4 in the presence of ADP and magnesium. Probably acts as the rate-limiting enzyme of the InsP6 pathway. Modifies TNF-alpha-induced apoptosis

by interfering with the activation of TNFRSF1A-associated death domain.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Immunohistochemistry of paraffin-embedded human liver tissue using [TA386860] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using [TA386860] at dilution of 1:100





Western blot

All lanes: ITPK1 antibody at 3.58µg/ml

Lane 1: Hela whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 46, 36 kDa Observed band size: 46 kDa