

## Product datasheet for **TA386865**


### DDX19A Rabbit Polyclonal Antibody

#### Product data:


|                       |  |
|-----------------------|--|
| Product Type:         | Primary Antibodies   |
| Applications:         | IHC, WB  |
| Recommended Dilution: | Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200   |
| Reactivity:           | Mouse, Human   |
| Host:                 | Rabbit   |
| Isotype:              | IgG  |
| Clonality:            | Polyclonal   |
| Immunogen:            | Recombinant Human ATP-dependent RNA helicase DDX19A protein (1-478AA)  |
| Formulation:          | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4   |
| Concentration:        | lot specific   |
| Purification:         | >95%, Protein G purified   |
| Conjugation:          | Unconjugated   |
| Storage:              | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| Stability:            | 1 year from dispatch.  |
| Database Link:        | <a href="#">Q9NUU7</a>   |
| Background:           | ATP-dependent RNA helicase involved in mRNA export from the nucleus. Rather than unwinding RNA duplexes, DDX19 functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding proteins |




[View online »](#)

**Product images:**

Immunohistochemistry of paraffin-embedded human ovarian cancer using TA386865 at dilution of 1:100



Immunohistochemistry of paraffin-embedded human prostate cancer using TA386865 at dilution of 1:100



Western blot

All lanes: DDX19A antibody at 2 $\mu$ g/ml + Mouse gonad tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 54, 44 kDa

Observed band size: 54 kDa