

## **Product datasheet for TA387000**

## Product data:

**Product Type:** Primary Antibodies

**EIF3A Rabbit Polyclonal Antibody** 

Applications: IF, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Eukaryotic translation initiation factor 3 subunit A protein (1288-

1379AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

Conjugation: Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q14152

**Background:** Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required

for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Essential for the initiation of translation on type-1 viral ribosomal entry sites (IRESs), like for HCV, PV, EV71 or BEV translation (PubMed:23766293, PubMed:24357634). In case of FCV infection, plays a role in the ribosomal termination-reinitiation event leading to the

translation of VP2 (PubMed:18056426).




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunofluorescent analysis of PC-3 cells using TA387000 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blot

Positive WB detected in: Jurkat whole cell lysate

All lanes: EIF3A antibody at 4µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 167, 163 kDa Observed band size: 167 kDa