

## **Product datasheet for TA387111M**

## **HOOK1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200

Reactivity: Human Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Protein Hook homolog 1 protein (301-600AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9U|C3

**Background:** Required for spermatid differentiation. Probably involved in the positioning of the

microtubules of the manchette and the flagellum in relation to the membrane skeleton. Component of the FTS/Hook/FHIP complex (FHF complex). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex

(the HOPS complex).




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded human testis tissue using [TA387111] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human colon cancer using [TA387111] at dilution of 1:100





Western blot
All lanes: HOOK1 antibody at 4µg/ml
Lane 1: 293T whole cell lysate
Lane 2: MCF-7 whole cell lysate
Secondary
Goat polyclonal to rabbit IgG at 1/10000 dilution
Predicted band size: 85, 81 kDa
Observed band size: 85 kDa