

Product datasheet for TA387115

VANGL2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200

Reactivity: Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Vang-like protein 2 protein (1-110AA)

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Antigen Affinity Purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q9ULK5

Background: Involved in the control of early morphogenesis and patterning of both axial midline

structures and the development of neural plate. Plays a role in the regulation of planar cell polarity, particularly in the orientation of stereociliary bundles in the cochlea. Required for polarization and movement of myocardializing cells in the outflow tract and seems to act via RHOA signaling to regulate this process. Required for cell surface localization of FZD3 and

FZD6 in the inner ear (By similarity).




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:





Immunohistochemistry of paraffin-embedded human lung cancer using TA387115 at dilution of 1:100



Immunohistochemistry of paraffin-embedded human gastric cancer using TA387115 at dilution of 1:100







Western blot

All lanes: VANGL2 antibody at 2.26µg/ml

Lane 1: HepG2 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: Hela whole cell lysate Lane 4: NIH/3T3 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 60 kDa Observed band size: 60 kDa

Western Blot

Positive WB detected in: HepG2 whole cell lysate, MCF-7 whole cell lysate, Mouse kidney tissue

All lanes: Vangl2 antibody at 3μg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 60 kDa Observed band size: 60 kDa