

## **Product datasheet for TA387175M**

## **CHRDL1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Chordin-like protein 1 protein (92-237AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9BU40

**Background:** Antagonizes the function of BMP4 by binding to it and preventing its interaction with

receptors. Alters the fate commitment of neural stem cells from gliogenesis to neurogenesis. Contributes to neuronal differentiation of neural stem cells in the brain by preventing the adoption of a glial fate. May play a crucial role in dorsoventral axis formation. May play a role in embryonic bone formation (By similarity). May also play an important role in regulating retinal angiogenesis through modulation of BMP4 actions in endothelial cells. Plays a role

during anterior segment eye development.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



IHC image of [TA387175] diluted at 1:200 and staining in paraffin-embedded human endometrial cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot
Positive WB detected in: Hela whole cell lysate,
A549 whole cell lysate
All lanes: CHRDL1 antibody at 5.7µg/ml
Secondary
Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 52, 44, 53 kDa
Observed band size: 52 kDa