

## **Product datasheet for TA387177**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

# **CPNE7 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200

**Reactivity:** Mouse, Rat, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant Human Copine-7 protein (254-393AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

Conjugation: Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9UBL6


**Background:** Calcium-dependent phospholipid-binding protein that may play a role in calcium-mediated

intracellular processes.






### **Product images:**



Immunofluorescence staining of A549 cells with TA387177 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).




#### Western Blot

Positive WB detected in: U87 whole cell lysate, SH-SY5Y whole cell lysate, HepG2 whole cell lysate, Mouse brain tissue, Rat heart tissue All lanes: CPNE7 antibody at 3.6µg/ml Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 71, 62 kDa Observed band size: 62 kDa





IHC image of TA387177 diluted at 1:300 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.