

Product datasheet for TA387251M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DAD1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit

DAD1 protein (1-113AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: P61803

Background: Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a

defined glycan (Glc3Man9GlcNAc2 in eukaryotes) from the lipid carrier dolichol-

pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent

polypeptide chains, the first step in protein N-glycosylation (PubMed:22467853). N-


glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity (By similarity). Required for the assembly of both SST3A- and SS3B-containing OST complexes.

Loss of the DAD1 protein triggers apoptosis (PubMed:22467853).







Product images:



Immunohistochemistry of paraffin-embedded human prostate tissue using [TA387251] at dilution of 1:100



Immunofluorescent analysis of HepG2 cells using [TA387251] at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western blot

All lanes: DAD1 antibody at 2µg/ml Lane 1: EC109 whole cell lysate Lane 2: 293T whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/15000 dilution

Predicted band size: 13 kDa Observed band size: 13 kDa