

Product datasheet for TA387272

FOXE3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200

Reactivity: Rat, Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Forkhead box protein E3 protein (4-74AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q13461

Background: Transcription factor that controls lens epithelial cell growth through regulation of

proliferation, apoptosis and cell cycle (PubMed:22527307, PubMed:25504734). During lens

development, controls the ratio of the lens fiber cells to the cells of the anterior lens

epithelium by regulating the rate of proliferation and differentiation (By similarity). Controls

lens vesicle closure and subsequent separation of the lens vesicle from ectoderm (By similarity). Controls the expression of DNAJB1 in a pathway that is crucial for the

development of the anterior segment of the eye (PubMed:27218149).




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Immunofluorescence staining of HepG2 cells with TA387272 at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western Blot

Positive WB detected in: Mouse brain tissue, Rat

brain tissue

All lanes: FOXE3 antibody at 4µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 34 kDa Observed band size: 34 kDa