

Product datasheet for TA387329M

PEX13 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500

Reactivity: Rat, Human

Rabbit Host: Isotype: lgG

Polyclonal Clonality:

Immunogen: Recombinant Human Peroxisomal membrane protein PEX13 protein (1-130AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q92968

Background: Component of the peroxisomal translocation machinery with PEX14 and PEX17. Functions as

a docking factor for the predominantly cytoplasmic PTS1 receptor (PAS10/PEX5). Involved in

the import of PTS1 and PTS2 proteins.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



IHC image of [TA387329] diluted at 1:200 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot Positive WB detected in: Rat brain tissue All lanes: PEX13 antibody at 3.4µg/ml Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 45 kDa

Observed band size: 45 kDa