

## **Product datasheet for TA387414M**

## MTIF3 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200

**Reactivity:** Mouse, Rat, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Translation initiation factor IF-3, mitochondrial protein (1-278AA)

**Formulation:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9H2K0

**Background:** IF-3 binds to the 28S ribosomal subunit and shifts the equilibrum between 55S ribosomes and

their 39S and 28S subunits in favor of the free subunits, thus enhancing the availability of 28S

subunits on which protein synthesis initiation begins.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded human prostate cancer using [TA387414] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human lung cancer using [TA387414] at dilution of 1:100





Western blot

All lanes: MTIF3 antibody at 1.15 μg/ml Lane 1: Mouse small intestine tissue

Lane 2: Rat heart tissue Lane 3: Mouse liver tissue Lane 4: Hela whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 32 kDa Observed band size: 32, 100, 75 kDa