

## **Product datasheet for TA387443M**

## **ADAMTS4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:500

**Reactivity:** Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant Human A disintegrin and metalloproteinase with thrombospondin motifs 4

protein (213-319AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

Conjugation: Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: 075173

**Background:** Cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover. May play an

important role in the destruction of aggrecan in arthritic diseases. Could also be a critical factor in the exacerbation of neurodegeneration in Alzheimer disease. Cleaves aggrecan at

the '392-Glu-|-Ala-393' site.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Western Blot
Positive WB detected in: Mouse kidney tissue
All lanes: ADAMTS4 antibody at 3.2µg/ml
Secondary
Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 91 kDa
Observed band size: 91 kDa




Immunohistochemistry of paraffin-embedded human heart tissue using [TA387443] at dilution of 1:100





Immunohistochemistry of paraffin-embedded human liver tissue using [TA387443] at dilution of 1:100



IHC image of [TA387443] diluted at 1:250 and staining in paraffin-embedded human lung tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.