

## **Product datasheet for TA387450**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **C1D Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Recommended dilution: WB:1:2000-1:10000, IHC:1:50-1:500, IF:1:500-1:1000

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Nuclear nucleic acid-binding protein C1D protein (1-135AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q13901

**Background:** Plays a role in the recruitment of the RNA exosome complex to pre-rRNA to mediate the 3'-5'


end processing of the 5.8S rRNA; this function may include MPHOSPH6. Can activate PRKDC not only in the presence of linear DNA but also in the presence of supercoiled DNA. Can induce apoptosis in a p53/TP53 dependent manner. May regulate the TRAX/TSN complex

formation. Potentiates transcriptional repression by NR1D1 and THRB.






## **Product images:**




Immunohistochemistry of paraffin-embedded human liver tissue using TA387450 at dilution of 1:100




Immunofluorescence staining of Hela cells with TA387450 at 1:533, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).





Immunohistochemistry of paraffin-embedded human thyroid tissue using TA387450 at dilution of 1:100



Western Blot Positive WB detected in: Hela whole cell lysate All lanes: C1D antibody at 3.2µg/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 17 kDa Observed band size: 17 kDa