

Product datasheet for TA387543

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

H2AC11 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ChIP, IF, IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Histone H2A type 1 protein (2-130AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: POCOS8

Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin,


limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication

and chromosomal stability. DNA accessibility is regulated via a complex set of post-


translational modifications of histones, also called histone code, and nucleosome remodeling.



Product images:




Immunohistochemistry of paraffin-embedded human lung cancer using TA387543 at dilution of 1:100




Chromatin Immunoprecipitation Hela (1.2*10⁶) were cross-linked with formaldehyde, sonicated, and immunoprecipitated with 4µg anti-HIST1H2AG or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers (CSB-PP010389HU) against the PGK1 promoter.






Immunofluorescent analysis of Hela cells using TA387543 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Immunohistochemistry of paraffin-embedded human testis tissue using TA387543 at dilution of 1:100



Western blot All lanes: HIST1H2AG antibody at $4\mu g/ml + HL60$ whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 15 kDa

Observed band size: 15 kDa