

## Product datasheet for **TA387631M**

### **H2BC4 Rabbit Polyclonal Antibody**


#### **Product data:**

Product Type:	Primary Antibodies
Applications:	ChIP, WB
Recommended Dilution:	Recommended dilution: WB:1:100-1:1000
Reactivity:	Mouse, Rat, Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide sequence around site of Ser (14) derived from Human Histone H2B type 1-C/E/F/G/I
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Concentration:	lot specific
Purification:	Antigen Affinity Purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	<u><a href="#">P62807</a></u>
Background:	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.




[View online »](#)

## Product images:



Chromatin Immunoprecipitation HeLa ( $4 \times 10^6$ ) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with  $5 \mu\text{g}$  anti-HIST1H2BC ([TA387631]) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the  $\beta$ -Globin promoter.



### Western Blot

Positive WB detected in: HeLa whole cell lysate, NIH/3T3 whole cell lysate, K562 whole cell lysate, A549 whole cell lysate, Rat kidney tissue

All lanes: HIST1H2BC antibody at  $0.9 \mu\text{g/ml}$

Secondary

Goat polyclonal to rabbit IgG at  $1/50000$  dilution

Predicted band size: 14 kDa

Observed band size: 14 kDa