

Product datasheet for TA387651M

Product datasireet for TAS6705 IN

H2BC4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC, WB

Recommended Dilution: Recommended dilution: WB:1:100-1:1000, ICC:1:20-1:200

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide sequence around site of β-hydroxybutyryl-Lys (20) derived from Human Histone H2B

type 1-C/E/F/G/I

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: Antigen Affinity Purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: P62807

Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin,

limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication

and chromosomal stability. DNA accessibility is regulated via a complex set of post-

translational modifications of histones, also called histone code, and nucleosome remodeling.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Immunocytochemistry analysis of [TA387651] diluted at 1:40 and staining in Hela cells (treated with 50mM sodium 3-hydroxybutyrate for 72h) performed on a Leica BondTM system. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot

Detected samples: 293 whole cell lysate, A549 whole cell lysate; Untreated (-) or treated (+) with 50mM Sodium 3-hydroxybutyrate for 72h All lanes: HIST1H2BC antibody at 1:100 Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 14 kDa Observed band size: 14 kDa